



# Samoa Meteorology Division Early Action Rainfall Watch (EAR Watch)



The EAR Watch provides disaster managers' with a brief summary of recent rainfall patterns, particularly drought and the rainfall outlook for the coming months. Contact the Samoa Meteorology Division for further climate information.


**Issued: February 2019**

## Rainfall Status:

**Meteorological Drought** exists at Faleolo at the 6-month timescales while a **Drought Watch** is current for Nafanua at the 6 month timescales. **Drought Warning** are current at Apia for 3 and 6 months scale as well as Nafanua and Faleolo at 3 months scale.

## Rainfall Outlook:

**Alert 1 Dry** is anticipated for Apia while **Alert 1 Wet** is expected for the other three stations.




Currently, the ENSO is neutral with 50% chance of El Niño to occur in mid-2019.

*Note: Colours do not relate to those below.*

## Rainfall Status at 31 January 2019 and Outlook for March to May 2019

Rainfall Stations*	Rainfall Status			Rainfall Outlook
	Past 12 months	Past 6 months	Past 3 months	Next 3 months
Apia	Normal	Warning	Warning	Alert 1 Dry
Afiamalu	Normal	Normal	Normal	Alert 1 Wet
Nafanua	Normal	Watch	Warning	Alert 1 Wet
Faleolo	Normal	Drought	Warning	Alert 1 Wet



← Increasing chance of drier 3 months

Increasing chance of wetter 3 months →

## Rainfall Status

The World Meteorological Organization (WMO) recognised Percentile method has been used to assess rainfall status. Meteorological Drought is defined as drought assessed by historical rainfall data only.

## Rainfall Outlook

Seasonal outlooks have been produced using SCOPIC v4.4.16 which is a decision-support tool used to generate outlooks for temperature, rainfall and other climate related factors

<http://cosppac.bom.gov.au/products-and-services/seasonal-climate-outlooks-in-pacific-island-countries/>.

The outlook provides an indication of total three-month rainfall, not how intense the rain may be in any one event, nor how it may vary within the three months. A station is assigned 'No Alert' when near normal rainfall is favoured or there are equal chances of below normal, normal and above normal rainfall. Two months before the dry and wet season, until the start of the season, a second outlook is presented for the upcoming dry or wet season.


### Time periods and impacts

The following table provides examples of impacts that have been associated with drought at the 3, 6 and 12 month periods. Note the periods are estimates only and impacts overlap. Allow for uncertainty associated with seasonality, island size, topography, geology, soil type or socio-economic circumstances. Contact the National Disaster Management Office and relevant Department for further information on impacts.

Sector/ Department	12-month period is most relevant for	6-month period is most relevant for	3-month period is most relevant for
<b>Water</b>	major rivers, deep bores/large aquifer system, reservoirs, dams	small rivers, shallow bores, reservoirs	rainwater tanks, streams, shallow bores
<b>Environment/ Agriculture</b>	coconuts, breadfruit, mango, banana, fruit trees (nonu, lemon, orange), root crops (yam, taamu, cassava), kava	corn, pineapple, pawpaw, taro, kumala, avocado, cocoa, coffee	traditional vegetables, cabbage, tomatoes, beans, eggplant, watermelon, pasture
<b>Fire</b>	All Fires	Structural Fire	Bush and Rubbish Fire
<b>Health</b>	National public health impacts	Increasing public health impacts	Isolated public health impacts
<b>Tourism</b>			

### Samoa rainfall monitoring stations

\* Only stations over 25 complete years of rainfall data are used in the EAR Watch.



Contact the Samoa Meteorology Division for further information.

The Director, Samoa Meteorology Division  
P.O. Box 3020, Apia, Samoa,  
Phone: 685 20855 / 20856 Fax: 685 20857

Website: <http://www.samet.gov.ws/> Email: [ausetalia.titimaea@mnre.gov.ws](mailto:ausetalia.titimaea@mnre.gov.ws)